

Hindawi Publishing Corporation
Journal of Inequalities and Applications
Volume 2009, Article ID 851360, 10 pages
doi:10.1155/2009/851360

Research Article

Some New Hilbert's Type Inequalities

Chang-Jian Zhao¹ and Wing-Sum Cheung²

¹ Department of Information and Mathematics Sciences, College of Science,
China Jiliang University, Hangzhou 310018, China

² Department of Mathematics, The University of Hong Kong, Pokfulam Road, Hong Kong

Correspondence should be addressed to Chang-Jian Zhao, chjzhao@163.com

Received 25 December 2008; Accepted 24 April 2009

Recommended by Peter Pang

Some new inequalities similar to Hilbert's type inequality involving series of nonnegative terms are established.

Copyright © 2009 C.-J. Zhao and W.-S. Cheung. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In recent years, several authors [1–10] have given considerable attention to Hilbert's type inequalities and their various generalizations. In particular, in [1], Pachpatte proved some new inequalities similar to Hilbert's inequality [11, page 226] involving series of nonnegative terms. The main purpose of this paper is to establish their general forms.

2. Main Results

In [1], Pachpatte established the following inequality involving series of nonnegative terms.

Theorem A. Let $p \geq 1, q \geq 1$, and let $\{a_m\}$ and $\{b_n\}$ be two nonnegative sequences of real numbers defined for $m = 1, \dots, k$, and $n = 1, \dots, r$, where k, r are natural numbers. Let $A_m = \sum_{s=1}^m a_s$ and $B_n = \sum_{t=1}^n b_t$. Then

$$\sum_{m=1}^k \sum_{n=1}^r \frac{A_m^p B_n^q}{m+n} \leq C(p, q, k, r) \left(\sum_{m=1}^k (k-m+1) (a_m A_m^{p-1})^2 \right)^{1/2} \times \left(\sum_{n=1}^r (r-n+1) (b_n B_n^{q-1})^2 \right)^{1/2}, \quad (2.1)$$

where

$$C(p, q, k, r) = \frac{1}{2}pq(kr)^{1/2}. \quad (2.2)$$

We first establish the following general form of inequality (2.1).

Theorem 2.1. Let $p \geq 1, q \geq 1, t > 0$, and $1/\alpha + 1/\beta = 1, \alpha > 1$. Let $\{a_{m_1, \dots, m_n}\}$, and $\{b_{n_1, \dots, n_n}\}$ be positive sequences of real numbers defined for $m_i = 1, 2, \dots, k_i$, and $n_i = 1, 2, \dots, r_i$, where k_i, r_i ($i = 1, \dots, n$) are natural numbers. Let $A_{m_1, \dots, m_n} = \sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} a_{s_1, \dots, s_n}$, and $B_{n_1, \dots, n_n} = \sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} b_{t_1, \dots, t_n}$. Then

$$\begin{aligned} & \sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \frac{\alpha \beta t^{1/\beta} A_{m_1, \dots, m_n}^p B_{n_1, \dots, n_n}^q}{m_1 \cdots m_n \beta + n_1 \cdots n_n \alpha t} \\ & \leq L(k_1, \dots, k_n, r_1, \dots, r_n, p, q, \alpha, \beta) \\ & \quad \times \left(\sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \prod_{j=1}^n (k_j - m_j + 1) (a_{m_1, \dots, m_n} A_{m_1, \dots, m_n}^{p-1})^\beta \right)^{1/\beta} \\ & \quad \times \left(\sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \prod_{j=1}^n (r_j - n_j + 1) (b_{n_1, \dots, n_n} B_{n_1, \dots, n_n}^{q-1})^\alpha \right)^{1/\alpha}, \end{aligned} \quad (2.3)$$

where

$$L(k_1, \dots, k_n, r_1, \dots, r_n, p, q, \alpha, \beta) = pq(k_1 \cdots k_n)^{1/\alpha} (r_1 \cdots r_n)^{1/\beta}. \quad (2.4)$$

Proof. By using the following inequality (see [12]):

$$\left(\sum_{m_1=1}^{n_1} \cdots \sum_{m_n=1}^{n_n} z_{m_1, \dots, m_n} \right)^p \leq p \sum_{m_1=1}^{n_1} \cdots \sum_{m_n=1}^{n_n} z_{m_1, \dots, m_n} \left(\sum_{k_1=1}^{m_1} \cdots \sum_{k_n=1}^{m_n} z_{k_1, \dots, k_n} \right)^{p-1}, \quad (2.5)$$

where $p \geq 1$ is a constant, and $z_{m_1, \dots, m_n} \geq 0$, $m_i = 1, 2, \dots, k_i$, $i = 1, 2, \dots, n$, we obtain

$$A_{m_1, \dots, m_n}^p \leq p \sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} a_{s_1, \dots, s_n} A_{s_1, \dots, s_n}^{p-1}. \quad (2.6)$$

Similarly, we have

$$B_{n_1, \dots, n_n}^q \leq q \sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} b_{t_1, \dots, t_n} B_{t_1, \dots, t_n}^{q-1}. \quad (2.7)$$

From (2.6) and (2.7), using Hölder's inequality [13] and the elementary inequality:

$$a^{1/\alpha} b^{1/\beta} \leq \frac{a}{\alpha t^{1/\beta}} + \frac{t^{1/\alpha} b}{\beta}, \quad (2.8)$$

where $1/\alpha + 1/\beta = 1$, $\alpha > 1$, $b > 0$, $a > 0$, and $t > 0$, we have

$$\begin{aligned} A_{m_1, \dots, m_n}^p B_{n_1, \dots, n_n}^q &\leq pq \left(\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} a_{s_1, \dots, s_n} A_{s_1, \dots, s_n}^{p-1} \right) \left(\sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} b_{t_1, \dots, t_n} B_{t_1, \dots, t_n}^{q-1} \right) \\ &\leq pq (m_1, \dots, m_n)^{1/\alpha} \left(\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} (a_{s_1, \dots, s_n} A_{s_1, \dots, s_n}^{p-1})^\beta \right)^{1/\beta} \\ &\quad \times (n_1, \dots, n_n)^{1/\beta} \left(\sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} (b_{t_1, \dots, t_n} B_{t_1, \dots, t_n}^{q-1})^\alpha \right)^{1/\alpha} \\ &\leq pq \left(\frac{m_1 \cdots m_n}{\alpha t^{1/\beta}} + \frac{n_1 \cdots n_n t^{1/\alpha}}{\beta} \right) \left(\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} (a_{s_1, \dots, s_n} A_{s_1, \dots, s_n}^{p-1})^\beta \right)^{1/\beta} \\ &\quad \times \left(\sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} (b_{t_1, \dots, t_n} B_{t_1, \dots, t_n}^{q-1})^\alpha \right)^{1/\alpha}. \end{aligned} \quad (2.9)$$

Dividing both sides of (2.9) by $((m_1 \cdots m_n \beta + n_1 \cdots n_n \alpha t) / \alpha \beta t^{1/\beta})$, summing up over n_i from 1 to r_i ($i = 1, 2, \dots, n$) first, then summing up over m_i from 1 to k_i ($i = 1, 2, \dots, n$), using again Hölder's inequality, then interchanging the order of summation, we obtain

$$\begin{aligned} &\sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \frac{\alpha \beta t^{1/\beta} A_{m_1, \dots, m_n}^p B_{n_1, \dots, n_n}^q}{m_1 \cdots m_n \beta + n_1 \cdots n_n \alpha t} \\ &\leq pq \left\{ \sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \left(\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} (a_{s_1, \dots, s_n} A_{s_1, \dots, s_n}^{p-1})^\beta \right)^{1/\beta} \right\} \\ &\quad \times \left\{ \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \left(\sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} (b_{t_1, \dots, t_n} B_{t_1, \dots, t_n}^{q-1})^\alpha \right)^{1/\alpha} \right\} \\ &\leq pq (k_1 \cdots k_n)^{1/\alpha} \left\{ \sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \left(\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} (a_{s_1, \dots, s_n} A_{s_1, \dots, s_n}^{p-1})^\beta \right) \right\}^{1/\beta} \\ &\quad \times (r_1 \cdots r_n)^{1/\beta} \left\{ \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \left(\sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} (b_{t_1, \dots, t_n} B_{t_1, \dots, t_n}^{q-1})^\alpha \right) \right\}^{1/\alpha} \end{aligned}$$

$$\begin{aligned}
&= L(k_1, \dots, k_n, r_1, \dots, r_n, p, q, \alpha, \beta) \\
&\quad \times \left\{ \sum_{s_1=1}^{k_1} \cdots \sum_{s_n=1}^{k_n} (a_{s_1, \dots, s_n} A_{s_1, \dots, s_n}^{p-1})^\beta \left(\sum_{m_1=s_1}^{k_1} \cdots \sum_{m_n=s_n}^{k_n} 1 \right) \right\}^{1/\beta} \\
&\quad \times \left\{ \sum_{t_1=1}^{r_1} \cdots \sum_{t_n=1}^{r_n} (b_{t_1, \dots, t_n} B_{t_1, \dots, t_n}^{q-1})^\alpha \left(\sum_{n_1=t_1}^{r_1} \cdots \sum_{n_n=t_n}^{r_n} 1 \right) \right\}^{1/\alpha} \\
&= L(k_1, \dots, k_n, r_1, \dots, r_n, p, q, \alpha, \beta) \\
&\quad \times \left\{ \sum_{s_1=1}^{k_1} \cdots \sum_{s_n=1}^{k_n} \prod_{j=1}^n (k_j - s_j + 1) (a_{s_1, \dots, s_n} A_{s_1, \dots, s_n}^{p-1})^\beta \right\}^{1/\beta} \\
&\quad \times \left\{ \sum_{t_1=1}^{r_1} \cdots \sum_{t_n=1}^{r_n} \prod_{j=1}^n (r_j - t_j + 1) (b_{t_1, \dots, t_n} B_{t_1, \dots, t_n}^{q-1})^\alpha \right\}^{1/\alpha} \\
&= L(k_1, \dots, k_n, r_1, \dots, r_n, p, q, \alpha, \beta) \\
&\quad \times \left\{ \sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \prod_{j=1}^n (k_j - m_j + 1) (a_{m_1, \dots, m_n} A_{m_1, \dots, m_n}^{p-1})^\beta \right\}^{1/\beta} \\
&\quad \times \left\{ \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \prod_{j=1}^n (r_j - n_j + 1) (b_{n_1, \dots, n_n} B_{n_1, \dots, n_n}^{q-1})^\alpha \right\}^{1/\alpha}.
\end{aligned} \tag{2.10}$$

This completes the proof. \square

Remark 2.2. Taking $\alpha = \beta = n = j = 2$, (2.3) becomes

$$\begin{aligned}
&\sum_{m=1}^{k_1} \sum_{m_2=1}^{k_2} \left(\sum_{n_1=1}^{r_1} \sum_{n_2=1}^{r_2} \frac{A_{m_1, m_2}^p B_{n_1, n_2}^q}{m_1 m_2 t^{-1/2} + n_1 n_2 t^{1/2}} \right) \\
&\leq \frac{1}{2} p q \sqrt{k_1 k_2 r_1 r_2} \left(\sum_{m=1}^{k_1} \sum_{m_2=1}^{k_2} (k_1 - m_1 + 1)(k_2 - m_2 + 1) (a_{m_1, m_2} A_{m_1, m_2}^{p-1})^2 \right)^{1/2} \\
&\quad \times \left(\sum_{n=1}^{r_1} \sum_{n_2=1}^{r_2} (r_1 - n_1 + 1)(r_2 - n_2 + 1) (b_{n_1, n_2} B_{n_1, n_2}^{q-1})^2 \right)^{1/2}.
\end{aligned} \tag{2.11}$$

Taking $t = 1$, and changing $\{a_{m_1, m_2}\}$, $\{b_{n_1, n_2}\}$, $\{A_{m_1, m_2}\}$, and $\{B_{n_1, n_2}\}$ into $\{a_m\}$, $\{b_n\}$, $\{A_m\}$, and $\{B_n\}$, respectively, and with suitable changes, (2.11) reduces to Pachpatte [1, inequality (1)].

In [1], Pachpatte also established the following inequality involving series of nonnegative terms.

Theorem B. Let $\{a_m\}$, $\{b_n\}$, A_m, B_n be as defined in Theorem A. Let $\{p_m\}$ and $\{q_n\}$ be positive sequences for $m = 1, \dots, k$, and $n = 1, \dots, r$, where k, r are natural numbers. Define $P_m = \sum_{s=1}^m p_s$, and $Q_n = \sum_{t=1}^n q_t$. Let ϕ and ψ be real-valued, nonnegative, convex, submultiplicative functions defined on $\mathbb{R}_+ = [0, +\infty)$. Then

$$\begin{aligned} \sum_{m=1}^k \sum_{n=1}^r \frac{\phi(A_m)\psi(B_n)}{m+n} &\leq M(k, r) \left(\sum_{m=1}^k (k-m+1) \left(p_m \phi\left(\frac{a_m}{p_m}\right) \right)^2 \right)^{1/2} \\ &\quad \times \left(\sum_{n=1}^r (r-n+1) \left(q_n \psi\left(\frac{b_n}{q_n}\right) \right)^2 \right)^{1/2}, \end{aligned} \quad (2.12)$$

where

$$M(k, r) = \frac{1}{2} \left(\sum_{m=1}^k \left(\frac{\phi(P_m)}{P_m} \right)^2 \right)^{1/2} \left(\sum_{n=1}^r \left(\frac{\psi(Q_n)}{Q_n} \right)^2 \right)^{1/2}. \quad (2.13)$$

Inequality (2.12) can also be generalized to the following general form.

Theorem 2.3. Let $\{a_{m_1, \dots, m_n}\}$, $\{b_{n_1, \dots, n_n}\}$, $\alpha, \beta, t, A_{m_1, \dots, m_n}$, and B_{n_1, \dots, n_n} be as defined in Theorem 2.1. Let $\{p_{m_1, \dots, m_n}\}$ and $\{q_{n_1, \dots, n_n}\}$ be positive sequences for $m_i = 1, 2, \dots, k_i$, and $n_i = 1, 2, \dots, r_i$ ($i = 1, 2, \dots, n$). Define $P_{m_1, \dots, m_n} = \sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} p_{s_1, \dots, s_n}$, and $Q_{n_1, \dots, n_n} = \sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} q_{t_1, \dots, t_n}$. Let ϕ and ψ be real-valued, nonnegative, convex, submultiplicative functions defined on $\mathbb{R}_+ = [0, +\infty)$. Then

$$\begin{aligned} \sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \frac{\alpha \beta t^{1/\beta} \phi(A_{m_1, \dots, m_n}) \psi(B_{n_1, \dots, n_n})}{m_1 \cdots m_n \beta + n_1 \cdots n_n \alpha t} \\ \leq M(k_1, \dots, k_n, r_1, \dots, r_n, \alpha, \beta) \\ \times \left\{ \sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \prod_{j=1}^n (k_j - m_j + 1) \left(p_{m_1, \dots, m_n} \phi\left(\frac{a_{m_1, \dots, m_n}}{p_{m_1, \dots, m_n}}\right) \right)^\beta \right\}^{1/\beta} \\ \times \left\{ \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \prod_{j=1}^n (r_j - n_j + 1) \left(q_{n_1, \dots, n_n} \psi\left(\frac{b_{n_1, \dots, n_n}}{q_{n_1, \dots, n_n}}\right) \right)^\alpha \right\}^{1/\alpha}, \end{aligned} \quad (2.14)$$

where

$$\begin{aligned} M(k_1, \dots, k_n, r_1, \dots, r_n, \alpha, \beta) \\ = \left(\sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \left(\frac{\phi(P_{m_1, \dots, m_n})}{P_{m_1, \dots, m_n}} \right)^\alpha \right)^{1/\alpha} \left(\sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \left(\frac{\psi(Q_{n_1, \dots, n_n})}{Q_{n_1, \dots, n_n}} \right)^\beta \right)^{1/\beta}. \end{aligned} \quad (2.15)$$

Proof. By the hypotheses, Jensen's inequality, and Hölder's inequality, we obtain

$$\begin{aligned}
 \phi(A_{m_1, \dots, m_n}) &= \phi\left(\frac{P_{m_1, \dots, m_n} \sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} p_{s_1, \dots, s_n} (a_{s_1, \dots, s_n} / p_{s_1, \dots, s_n})}{\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} p_{s_1, \dots, s_n}}\right) \\
 &\leq \phi(P_{m_1, \dots, m_n}) \phi\left(\frac{\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} p_{s_1, \dots, s_n} (a_{s_1, \dots, s_n} / p_{s_1, \dots, s_n})}{\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} p_{s_1, \dots, s_n}}\right) \\
 &\leq \frac{\phi(P_{m_1, \dots, m_n})}{P_{m_1, \dots, m_n}} \sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} p_{s_1, \dots, s_n} \phi\left(\frac{a_{s_1, \dots, s_n}}{p_{s_1, \dots, s_n}}\right) \\
 &\leq \frac{\phi(P_{m_1, \dots, m_n})}{P_{m_1, \dots, m_n}} (m_1 \cdots m_n)^{1/\alpha} \left(\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} \left(p_{s_1, \dots, s_n} \phi\left(\frac{a_{s_1, \dots, s_n}}{p_{s_1, \dots, s_n}}\right)\right)^\beta\right)^{1/\beta}.
 \end{aligned} \tag{2.16}$$

Similarly,

$$\phi(B_{n_1, \dots, n_n}) \leq \frac{\phi(Q_{n_1, \dots, n_n})}{Q_{n_1, \dots, n_n}} (n_1 \cdots n_n)^{1/\beta} \left(\sum_{n_1=1}^{n_1} \cdots \sum_{n_n=1}^{n_n} \left(q_{t_1, \dots, t_n} \psi\left(\frac{b_{t_1, \dots, t_n}}{q_{t_1, \dots, t_n}}\right)\right)^\alpha\right)^{1/\alpha}. \tag{2.17}$$

By (2.16) and (2.17), and using the elementary inequality:

$$a^{1/\alpha} b^{1/\beta} \leq \frac{a}{\alpha t^{1/\beta}} + \frac{t^{1/\alpha} b}{\beta}, \tag{2.18}$$

where $1/\alpha + 1/\beta = 1$, $\alpha > 1$, $b > 0$, $a > 0$, and $t > 0$, we have

$$\begin{aligned}
 \phi(A_{m_1, \dots, m_n}) \phi(B_{n_1, \dots, n_n}) &\leq \left(\frac{m_1 \cdots m_n}{\alpha t^{1/\beta}} + \frac{n_1 \cdots n_n t^{1/\alpha}}{\beta}\right) \\
 &\quad \times \frac{\phi(P_{m_1, \dots, m_n})}{P_{m_1, \dots, m_n}} \left(\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} \left(p_{s_1, \dots, s_n} \phi\left(\frac{a_{s_1, \dots, s_n}}{p_{s_1, \dots, s_n}}\right)\right)^\beta\right)^{1/\beta} \\
 &\quad \times \frac{\phi(Q_{n_1, \dots, n_n})}{Q_{n_1, \dots, n_n}} \left(\sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} \left(q_{t_1, \dots, t_n} \psi\left(\frac{b_{t_1, \dots, t_n}}{q_{t_1, \dots, t_n}}\right)\right)^\alpha\right)^{1/\alpha}.
 \end{aligned} \tag{2.19}$$

Dividing both sides of (2.19) by $(m_1 \cdots m_n \beta + n_1 \cdots n_n \alpha t / \alpha \beta t^{1/\beta})$, and summing up over n_i from 1 to r_i ($i = 1, 2, \dots, n$) first, then summing up over m_i from 1 to k_i ($i = 1, 2, \dots, n$), using again inverse Hölder's inequality, and then interchanging the order of summation, we obtain

$$\begin{aligned}
& \sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \frac{\alpha \beta t^{1/\beta} \phi(A_{m_1, \dots, m_n}) \psi(B_{n_1, \dots, n_n})}{m_1 \cdots m_n \beta + n_1 \cdots n_n \alpha t} \\
& \leq \sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \left(\frac{\phi(P_{m_1, \dots, m_n})}{P_{m_1, \dots, m_n}} \left(\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} \left(p_{s_1, \dots, s_n} \phi \left(\frac{a_{s_1, \dots, s_n}}{p_{s_1, \dots, s_n}} \right) \right)^\beta \right)^{1/\beta} \right) \\
& \quad \times \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \left(\frac{\phi(Q_{n_1, \dots, n_n})}{Q_{n_1, \dots, n_n}} \left(\sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} \left(q_{t_1, \dots, t_n} \psi \left(\frac{b_{t_1, \dots, t_n}}{q_{t_1, \dots, t_n}} \right) \right)^\alpha \right)^{1/\alpha} \right) \\
& \leq \left(\sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \left(\frac{\phi(P_{m_1, \dots, m_n})}{P_{m_1, \dots, m_n}} \right)^\alpha \right)^{1/\alpha} \\
& \quad \times \left(\sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \left(\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} \left(p_{s_1, \dots, s_n} \phi \left(\frac{a_{s_1, \dots, s_n}}{p_{s_1, \dots, s_n}} \right) \right)^\beta \right) \right)^{1/\beta} \\
& \quad \times \left(\sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \left(\frac{\psi(Q_{n_1, \dots, n_n})}{Q_{n_1, \dots, n_n}} \right)^\beta \right)^{1/\beta} \\
& \quad \times \left(\sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \left(\sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} \left(q_{t_1, \dots, t_n} \psi \left(\frac{b_{t_1, \dots, t_n}}{q_{t_1, \dots, t_n}} \right) \right)^\alpha \right) \right)^{1/\alpha} \\
& = M(k_1, \dots, k_n, r_1, \dots, r_n, \alpha, \beta) \\
& \quad \times \left\{ \sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \prod_{j=1}^n (k_j - m_j + 1) \left(p_{m_1, \dots, m_n} \phi \left(\frac{a_{m_1, \dots, m_n}}{p_{m_1, \dots, m_n}} \right) \right)^\beta \right\}^{1/\beta} \\
& \quad \times \left\{ \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \prod_{j=1}^n (r_j - n_j + 1) \left(q_{n_1, \dots, n_n} \psi \left(\frac{b_{n_1, \dots, n_n}}{q_{n_1, \dots, n_n}} \right) \right)^\alpha \right\}^{1/\alpha}.
\end{aligned} \tag{2.20}$$

The proof is complete. \square

Remark 2.4. Taking $\alpha = \beta = n = j = 2$, (2.14) becomes

$$\begin{aligned} & \sum_{m=1}^{k_1} \sum_{m_2=1}^{k_2} \left(\sum_{n_1=1}^{r_1} \sum_{n_2=1}^{r_2} \frac{\phi(A_{m_1, m_2}) \psi(B_{n_1, n_2})}{m_1 m_2 t^{-1/2} + n_1 n_2 t^{1/2}} \right) \\ & \leq \overline{M}(k_1, k_2, r_1, r_2) \\ & \quad \times \left(\sum_{m=1}^{k_1} \sum_{m_2=1}^{k_2} (k_1 - m_1 + 1)(k_2 - m_2 + 1) \left(p_{m_1, m_2} \phi \left(\frac{a_{m_1, \dots, m_n}}{p_{m_1, \dots, m_n}} \right) \right)^2 \right)^{1/2} \\ & \quad \times \left(\sum_{n=1}^{r_1} \sum_{n_2=1}^{r_2} (r_1 - n_1 + 1)(r_2 - n_2 + 1) \left(q_{n_1, n_2} \psi \left(\frac{b_{n_1, \dots, n_n}}{q_{n_1, \dots, n_n}} \right) \right)^2 \right)^{1/2}, \end{aligned} \quad (2.21)$$

where

$$\overline{M}(k_1, k_2, r_1, r_2) = \left(\sum_{m_1=1}^{k_1} \sum_{m_2=1}^{k_2} \left(\frac{\phi(P_{m_1, m_2})}{P_{m_1, m_2}} \right)^2 \right)^{1/2} \left(\sum_{n_1=1}^{r_1} \sum_{n_2=1}^{r_2} \left(\frac{\psi(Q_{n_1, n_2})}{Q_{n_1, n_2}} \right)^2 \right)^{1/2}. \quad (2.22)$$

Taking $t = 1$, and changing $\{a_{m_1, m_2}\}$, $\{b_{n_1, n_2}\}$, $\{A_{m_1, m_2}\}$, and $\{B_{n_1, n_2}\}$ into $\{a_m\}$, $\{b_n\}$, $\{A_m\}$, and $\{B_n\}$, respectively, and with suitable changes, (2.21) reduces to Pachpatte [1, Inequality (7)].

Theorem 2.5. Let $\{a_{m_1, \dots, m_n}\}$, $\{b_{n_1, \dots, n_n}\}$, $\{p_{m_1, \dots, m_n}\}$, $\{q_{n_1, \dots, n_n}\}$, P_{m_1, \dots, m_n} , and Q_{n_1, \dots, n_n} , α, β, t , be as defined in Theorem 2.3. Define

$$\begin{aligned} A_{m_1, \dots, m_n} &= \frac{1}{P_{m_1, \dots, m_n}} \sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} p_{s_1, \dots, s_n} a_{s_1, \dots, s_n}, \\ B_{n_1, \dots, n_2} &= \frac{1}{Q_{n_1, \dots, n_n}} \sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} q_{t_1, \dots, t_n} b_{t_1, \dots, t_n}, \end{aligned} \quad (2.23)$$

for $m_i = 1, 2, \dots, k_i$, and $n_i = 1, 2, \dots, r_i$ ($i = 1, 2, \dots, n$), where k_i, r_i ($i = 1, \dots, n$) are natural numbers. Let ϕ and ψ be real-valued, nonnegative, convex functions defined on $\mathbb{R}_+ = [0, +\infty)$. Then

$$\begin{aligned} & \sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \frac{\alpha \beta t^{1/\beta} P_{m_1, \dots, m_n} Q_{n_1, \dots, n_n} \phi(A_{m_1, \dots, m_n}) \psi(B_{n_1, \dots, n_n})}{m_1 \cdots m_n \beta + n_1 \cdots n_n \alpha t} \\ & = (k_1 \cdots k_n)^{1/\alpha} (r_1 \cdots r_n)^{1/\beta} \\ & \quad \times \left\{ \sum_{m_1=1}^{k_1} \cdots \sum_{m_n=1}^{k_n} \prod_{j=1}^n (k_j - m_j + 1) (p_{m_1, \dots, m_n} \phi(a_{m_1, \dots, m_n}))^\beta \right\}^{1/\beta} \\ & \quad \times \left\{ \sum_{n_1=1}^{r_1} \cdots \sum_{n_n=1}^{r_n} \prod_{j=1}^n (r_j - n_j + 1) (q_{n_1, \dots, n_n} \psi(b_{n_1, \dots, n_n}))^\alpha \right\}^{1/\alpha}. \end{aligned} \quad (2.24)$$

Proof. By the hypotheses, Jensen's inequality, and Hölder's inequality, it is easy to observe that

$$\begin{aligned}\phi(A_{m_1, \dots, m_n}) &= \phi\left(\frac{1}{P_{m_1, \dots, m_n}} \sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} p_{s_1, \dots, s_n} a_{s_1, \dots, s_n}\right) \\ &\leq \frac{1}{P_{m_1, \dots, m_n}} \sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} p_{s_1, \dots, s_n} \phi(a_{s_1, \dots, s_n}) \\ &\leq \frac{1}{P_{m_1, \dots, m_n}} (m_1 \cdots m_n)^{1/\alpha} \left(\sum_{s_1=1}^{m_1} \cdots \sum_{s_n=1}^{m_n} (p_{s_1, \dots, s_n} \phi(a_{s_1, \dots, s_n}))^\beta \right)^{1/\beta},\end{aligned}\quad (2.25)$$

$$\begin{aligned}\psi(B_{n_1, \dots, n_n}) &= \psi\left(\frac{1}{Q_{n_1, \dots, n_n}} \sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} q_{t_1, \dots, t_n} b_{t_1, \dots, t_n}\right) \\ &\leq \frac{1}{Q_{n_1, \dots, n_n}} \sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} q_{t_1, \dots, t_n} \psi(b_{t_1, \dots, t_n}) \\ &\leq \frac{1}{Q_{n_1, \dots, n_n}} (n_1 \cdots n_n)^{1/\beta} \left(\sum_{t_1=1}^{n_1} \cdots \sum_{t_n=1}^{n_n} (q_{t_1, \dots, t_n} \psi(b_{t_1, \dots, t_n}))^\alpha \right)^{1/\alpha}.\end{aligned}\quad (2.26)$$

Proceeding now much as in the proof of Theorems 2.1 and 2.3, and with suitable modifications, it is not hard to arrive at the desired inequality. The details are omitted here. \square

Remark 2.6. In the special case where $j = 1$, $t = 1$, $\alpha = \beta = 2$, and $n = 1$, Theorem 2.5 reduces to the following result.

Theorem C. Let $\{a_m\}, \{b_n\}, \{p_m\}, \{q_n\}, P_m, Q_n$ be as defined in Theorem B. Define $A_m = (1/P_m) \sum_{s=1}^m p_s a_s$, and $B_n = (1/Q_n) \sum_{t=1}^n q_t b_t$ for $m = 1, \dots, k$, and $n = 1, \dots, r$, where k, r are natural numbers. Let ϕ and ψ be real-valued, nonnegative, convex functions defined on $\mathbb{R}_+ = [0, +\infty)$. Then

$$\begin{aligned}\sum_{m=1}^k \sum_{n=1}^r \frac{P_m Q_n \phi(A_m) \psi(B_n)}{m+n} &\leq \frac{1}{2} (kr)^{1/2} \left(\sum_{m=1}^k (k-m+1) (p_m \phi(a_m))^2 \right)^{1/2} \\ &\quad \times \left(\sum_{n=1}^r (r-n+1) (q_n \psi(b_n))^2 \right)^{1/2}.\end{aligned}\quad (2.27)$$

This is the new inequality of Pachpatte in [1, Theorem 4].

Remark 2.7. Taking $j = 1$, $t = 1$, $p_m = 1$, $q_n = 1$, $\alpha = \beta = 2$, and $n = 1$ in Theorem 2.5, and in view of $P_m = m$, $Q_n = n$, we obtain the following theorem.

Theorem D. Let $\{a_m\}$, $\{b_n\}$ be as defined in Theorem A. Define $A_m = (1/m) \sum_{s=1}^m a_s$, and $B_n = (1/n) \sum_{t=1}^n b_t$, for $m = 1, \dots, k$, and $n = 1, \dots, r$, where k, r are natural numbers. Let ϕ and ψ be real-valued, nonnegative, convex functions defined on $R_+ = [0, +\infty)$. Then

$$\sum_{m=1}^k \sum_{n=1}^r \frac{mn}{m+n} \phi(A_m) \psi(B_n) \leq \frac{1}{2} (kr)^{1/2} \left(\sum_{m=1}^k (k-m+1) (\phi(a_m))^2 \right)^{1/2} \times \left(\sum_{n=1}^r (r-n+1) (\psi(b_n))^2 \right)^{1/2}. \quad (2.28)$$

This is the new inequality of Pachpatte in [1, Theorem 3].

Acknowledgments

Research is supported by Zhejiang Provincial Natural Science Foundation of China (Y605065), Foundation of the Education Department of Zhejiang Province of China (20050392). Research is partially supported by the Research Grants Council of the Hong Kong SAR, China (Project no. HKU7016/07P) and a HKU Seed Grant for Basic Research.

References

- [1] B. G. Pachpatte, "On some new inequalities similar to Hilbert's inequality," *Journal of Mathematical Analysis and Applications*, vol. 226, no. 1, pp. 166–179, 1998.
- [2] G. D. Handley, J. J. Koliha, and J. E. Pečarić, "New Hilbert-Pachpatte type integral inequalities," *Journal of Mathematical Analysis and Applications*, vol. 257, no. 1, pp. 238–250, 2001.
- [3] M. Gao and B. Yang, "On the extended Hilbert's inequality," *Proceedings of the American Mathematical Society*, vol. 126, no. 3, pp. 751–759, 1998.
- [4] K. Jichang, "On new extensions of Hilbert's integral inequality," *Journal of Mathematical Analysis and Applications*, vol. 235, no. 2, pp. 608–614, 1999.
- [5] B. Yang, "On new generalizations of Hilbert's inequality," *Journal of Mathematical Analysis and Applications*, vol. 248, no. 1, pp. 29–40, 2000.
- [6] W. Zhong and B. Yang, "On a multiple Hilbert-type integral inequality with the symmetric kernel," *Journal of Inequalities and Applications*, vol. 2007, Article ID 27962, 17 pages, 2007.
- [7] C.-J. Zhao, "Inverses of disperse and continuous Pachpatte's inequalities," *Acta Mathematica Sinica*, vol. 46, no. 6, pp. 1111–1116, 2003.
- [8] C.-J. Zhao, "Generalization on two new Hilbert type inequalities," *Journal of Mathematics*, vol. 20, no. 4, pp. 413–416, 2000.
- [9] C.-J. Zhao and L. Debnath, "Some new inverse type Hilbert integral inequalities," *Journal of Mathematical Analysis and Applications*, vol. 262, no. 1, pp. 411–418, 2001.
- [10] G. D. Handley, J. J. Koliha, and J. E. Pečarić, "A Hilbert type inequality," *Tamkang Journal of Mathematics*, vol. 31, no. 4, pp. 311–315, 2000.
- [11] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge University Press, Cambridge, UK, 1934.
- [12] J. Németh, "Generalizations of the Hardy-Littlewood inequality," *Acta Scientiarum Mathematicarum*, vol. 32, pp. 295–299, 1971.
- [13] E. F. Beckenbach and R. Bellman, *Inequalities*, Ergebnisse der Mathematik und ihrer Grenzgebiete, N. F., Band 30, Springer, Berlin, Germany, 1961.